arXiv:1012.4669v2 [astro-ph.CO] 25 Jul 2011

G-essence with Yukawa Interactions

I.Kulnazarov[1], K.Yerzhanov[1], O.Razina[1], Sh.Myrzakul[1], P.Tsyba[1], R.Myrzakulov[1,2*]

[1]*Eurasian International Center for Theoretical Physics,*
Eurasian National University, Astana 010008, Kazakhstan
[2]*Department of Physics, CSU Fresno, Fresno, CA 93740 USA*

Abstract

We study the g-essence model with Yukawa interactions between a scalar field ϕ and a Dirac field ψ. For the homogeneous, isotropic and flat Friedmann–Robertson–Walker universe filled with the such g-essence, the exact solution of the model is found. Moreover, we reconstruct the corresponding scalar and fermionic potentials which describe the coupled dynamics of the scalar and fermionic fields. It is shown that some particular g-essence models with Yukawa interactions correspond to the usual and generalized Chaplygin gas unified models of dark energy and dark matter. Also we present some scalar–fermionic Dirac–Born–Infeld models corresponding g-essence models with Yukawa interactions which again describe the unified dark energy–dark matter system.

1 Introduction

One of the most puzzling discovery of the last years in physics is the current acceleration of the universe [1]–[2]. An unknown energy component, dubbed as dark energy, is proposed to explain this acceleration. Dark energy almost equally distributes in the universe, and its pressure is negative. The simplest and most theoretically appealing candidate of dark energy is the cosmological constant that is the ΛCDM model. In this case, the equation of state parameter $\omega = -1$. Although the ΛCDM model is in general agreement with the current astronomical observations, but has some difficulties e.g. to reconcile the small observational value of dark energy density with estimates from quantum field theories. So although the ΛCDM model is the most obvious choice, but it suffers from coincidence problem and the fine-tuning problems. It is thus natural to pursue alternative possibilities to explain the mystery of dark energy. In order to explain the acceleration that is dark energy, many kinds of models have been proposed, such as quintessence, phantom, k-essence, tachyon, f-essence, Chaplygin gas and its generalizations, etc.

In the last years, the k-essence model has received much attention. It was originally proposed as a model for inflation [3], and then as a model for dark energy [4]–[7]. Since from it was proposed, k-essence was been studied intensively. It is still worth investigating in a systematic way the possible cosmological behavior of the k-essence. Quite recently, the so-called g-essence model has been proposed [13], which is a more generalized model than k-essence. In fact, the g-essence contains, as particular cases, two important models: k-essence and f-essence. Note that f-essence is the fermionic counterpart of k-essence.

To our knowledge, in the literature there are relatively few works on the dark energy models with fermionic fields. However, in the recent years several approaches were made to explain the

---

[*] The corresponding author. Email: rmyrzakulov@gmail.com; rmyrzakulov@csufresno.edu

accelerated expansion by choosing fermionic fields as the gravitational sources of energy (see e.g. refs. [10]–[36]). In particular, it was shown that the fermionic field plays very important role in: i) isotropization of initially anisotropic spacetime; ii) formation of singularity free cosmological solutions; iii) explaining late-time acceleration. In particle physics, the Yukawa interaction plays an important role. It has the form

$$U = -g\bar{\psi}\phi\psi. \tag{1.1}$$

It describes the interaction between a scalar field ϕ and a Dirac field ψ. Some properties of the Yukawa interaction (1.1) related with the gravitational field were considered in [41]–[45]. With the Yukawa interaction (1.1) is related the so-called Yukawa potential,

$$V(x) = -g^2x^{-1}e^{-mx} \tag{1.2}$$

which is negative, that is, the corresponding force is attractive. The relation between the Yukawa potential (1.2) and the accelerated expansion of the universe were studied by some authors (see e.g. [46]). In this paper, we focus on so-called g-essence model [13] which is some hybrid construction of k-essence and f-essence. If exactly, we will consider the g-essence with the Yukawa interaction (1.1). The formulation of the gravity-fermionic theory has been discussed in detail elsewhere [37]–[40]., so we will only present the result here.

This paper is organized as follows. In the following section, we briefly review g-essence. In Sec. 3, we introduce the g-essence model with the Yukawa interaction. In Sec. 4, we construct the solution of the particular g-essence model with the scalar–fermionic Yukawa interaction. The unified scalar–fermionic Chaplygin gas model of dark energy and dark matter from the g-essence model with the Yukawa interaction were constructed in section 5 and its extension for the generalized Chaplygin gas case in section 6. The scalar–fermionic Dirac–Born–Infeld (DBI) counterpart of the g-essence model with the Yukawa interaction (3.1) was constructed in section 7. Finally, we shall close with a few concluding remarks in Sec. 8. The metric signature used is $(+,-,-,-)$ and units have been chosen so that $8\pi G = c = \hbar = 1$.

2 Basics of g-essence

The action of g-essence has the form [13]

$$S = \int d^4x\sqrt{-g}[R + 2K(X, Y, \phi, \psi, \bar{\psi})], \tag{2.1}$$

where K is the g-essence Lagrangian and is some function of its arguments, ϕ is a scalar function, $\psi = (\psi_1, \psi_2, \psi_3, \psi_4)^T$ is a fermionic function and $\bar{\psi} = \psi^+\gamma^0$ is its adjoint function, the curvature scalar R. Here

$$X = 0.5g^{\mu\nu}\nabla_\mu\phi\nabla_\nu\phi, \quad Y = 0.5i[\bar{\psi}\Gamma^\mu D_\mu\psi - (D_\mu\bar{\psi})\Gamma^\mu\psi] \tag{2.2}$$

are the canonical kinetic terms for the scalar and fermionic fields, respectively. ∇_μ and D_μ are the covariant derivatives. The fermionic fields are treated here as classically commuting fields. The model (2.1) admits important two reductions: *k-essence* and *f-essence*. In this sense, it is the more general essence model and in [13] it was called *g-essence*. Note that to find the equations of motion we need the variations

$$\begin{aligned}
\delta\sqrt{-g} &= -0.5g_{\mu\nu}\sqrt{-g}\delta g^{\mu\nu}, && (2.3)\\
\delta R &= (R_{\mu\nu} + g_{\mu\nu}\Box - \nabla_\mu\nabla_\nu)\delta g^{\mu\nu}, && (2.4)\\
\delta K &= K^{'}\delta\phi + K^{''}\delta\psi + K^{'''}\delta\bar{\psi}, && (2.5)
\end{aligned}$$

where $\nabla_\nu V_\mu \equiv \partial_\nu V_\mu - \Gamma^\sigma_{\mu\nu}V_\sigma$ is the covariant derivative of a vector V_μ and the curved d'Alembertian on a scalar ϕ is

$$\Box\phi = \frac{1}{\sqrt{-g}}\partial^\mu(\sqrt{-g}\partial_\mu\phi). \tag{2.6}$$

We now consider the dynamics of the homogeneous, isotropic and flat FRW universe filled with g-essence. In this case, the background line element reads

$$ds^2 = dt^2 - a^2(dx^2 + dy^2 + dz^2) \tag{2.7}$$

and the vierbein is chosen to be

$$(e_a^\mu) = diag(1, 1/a, 1/a, 1/a), \quad (e_\mu^a) = diag(1, a, a, a). \tag{2.8}$$

In the case of the FRW metric (2.7), the equations corresponding to the action (2.1) look like [13]

$$\begin{aligned}
3H^2 - \rho &= 0, \qquad &(2.9)\\
2\dot{H} + 3H^2 + p &= 0, &(2.10)\\
K_X\ddot{\phi} + (\dot{K}_X + 3HK_X)\dot{\phi} - K_\phi &= 0, &(2.11)\\
K_Y\dot{\psi} + 0.5(3HK_Y + \dot{K}_Y)\psi - i\gamma^0 K_{\bar{\psi}} &= 0, &(2.12)\\
K_Y\dot{\bar{\psi}} + 0.5(3HK_Y + \dot{K}_Y)\bar{\psi} + iK_\psi\gamma^0 &= 0, &(2.13)\\
\dot{\rho} + 3H(\rho + p) &= 0, &(2.14)
\end{aligned}$$

where $K_X = dK/dX, \quad K_Y = dK/dY, \quad K_\phi = dK/d\phi, \quad K_\psi = dK/d\psi, \quad K_{\bar{\psi}} = dK/d\bar{\psi}$ and $H = \dot{a}/a$ denotes the Hubble parameter, the dot represents a differentiation with respect to time t. Here the kinetic terms, the energy density and the pressure take the form

$$X = 0.5\dot{\phi}^2, \quad Y = 0.5i(\bar{\psi}\gamma^0\dot{\psi} - \dot{\bar{\psi}}\gamma^0\psi) \tag{2.15}$$

and

$$\rho = 2XK_X + YK_Y - K, \quad p = K. \tag{2.16}$$

Note that the equations of g-essence (2.9)-(2.14) can be rewritten as

$$\begin{aligned}
3H^2 - \rho &= 0, \qquad &(2.17)\\
2\dot{H} + 3H^2 + p &= 0, &(2.18)\\
(a^3K_X\dot{\phi})_t - a^3K_\phi &= 0, &(2.19)\\
(a^3K_Y\psi_j^2)_t - 2iK_{\bar{\psi}}(\gamma^0\psi)_j &= 0, &(2.20)\\
(a^3K_Y\psi_j^{*2})_t + 2iK_\psi(\bar{\psi}\gamma^0)_j &= 0, &(2.21)\\
\dot{\rho} + 3H(\rho + p) &= 0. &(2.22)
\end{aligned}$$

Also we present the useful formula

$$K_YY = 0.5iK_Y(\bar{\psi}\gamma^0\dot{\psi} - \dot{\bar{\psi}}\gamma^0\psi) = -0.5(K_\psi\psi + K_{\bar{\psi}}\bar{\psi}) \tag{2.23}$$

and the equation for $u = \bar{\psi}\psi$:

$$[\ln(ua^3K_Y)]_t u = -iK_Y^{-1}(\bar{\psi}\gamma^0K_{\bar{\psi}} - K_\psi\gamma^0\psi). \tag{2.24}$$

Finally, we note that some exact solutions of g-essence (2.9)–(2.14) were presented in [17]-[18] (see, also [14]-[16] in which some integrable aspects of the FRW models were considered).

2.1 Purely kinetic g-essence

Let us consider the purely kinetic g-essence, that is, when $K = K(X, Y)$. In this case, the system (2.9)–(2.14) becomes

$$\begin{aligned}
3H^2 - \rho &= 0, \qquad &(2.25)\\
2\dot{H} + 3H^2 + p &= 0, &(2.26)\\
a^3K_X\dot{\phi} - \sigma &= 0, &(2.27)\\
a^3K_Y\psi_j^2 - \varsigma_j &= 0, &(2.28)\\
a^3K_Y\psi_j^{*2} - \varsigma_j^* &= 0, &(2.29)\\
\dot{\rho} + 3H(\rho + p) &= 0, &(2.30)
\end{aligned}$$

where σ (ς) is the real (complex) constant. Hence we immediately get the solutions of the Klein–Gordon and Dirac equations, respectively, as

$$\phi = \sigma \int \frac{dt}{a^3 K_X}, \quad \psi_j = \sqrt{\frac{\varsigma_j}{a^3 K_Y}}. \tag{2.31}$$

Also the following useful formula holds:

$$X = \frac{0.5\sigma^2}{a^6 K_X^2} \quad or \quad K_X = \frac{\sigma}{a^3\sqrt{2X}}. \tag{2.32}$$

2.2 K-essence

Let us now we consider the following particular case of g-essence (2.1):

$$K = K_1 = K_1(X, \phi) \tag{2.33}$$

that corresponds to k-essence. Then the system (2.9)–(2.14) takes the form of the equations of k-essence (see e.g. [3]–[6])

$$3H^2 - \rho_k = 0, \tag{2.34}$$

$$2\dot{H} + 3H^2 + p_k = 0, \tag{2.35}$$

$$K_{1X}\ddot{\phi} + (\dot{K}_{1X} + 3HK_{1X})\dot{\phi} - K_{1\phi} = 0, \tag{2.36}$$

$$\dot{\rho}_k + 3H(\rho_k + p_k) = 0, \tag{2.37}$$

where the energy density and the pressure are given by

$$\rho_k = 2XK_{1X} - K_1, \quad p_k = K_1. \tag{2.38}$$

As is well known, the energy-momentum tensor for the k-essence field has the form

$$T_{\mu\nu} = K_X \nabla_\mu\phi\nabla_\nu\phi - g_{\mu\nu}K = 2K_X X u_{1\mu}u_{1\nu} - Kg_{\mu\nu} = (\rho_k + p_k)u_{1\mu}u_{1\nu} - p_k g_{\mu\nu}. \tag{2.39}$$

It is interesting to note that in the case of the FRW metric (2.7), purely kinetic k-essence and F(T)-gravity (modified teleparallel gravity) are equivalent to each other, if $a = e^{\pm\frac{\phi-\phi_0}{\sqrt{12}}}$ [8]–[9].

2.3 F-essence

Now we consider the following reduction of g-essence (2.1):

$$K = K_2 = K_2(Y, \psi, \bar{\psi}) \tag{2.40}$$

that is, f-essence [13]. The energy-momentum tensor for the f-essence field has the form

$$\begin{aligned} T_{\mu\nu} \equiv \frac{2}{\sqrt{-g}}\frac{\delta S}{\delta g_{\mu\nu}} &= 0.5iK_Y\left[\bar{\psi}\Gamma_{(\mu}D_{\nu)}\psi - D_{(\mu}\bar{\psi}\Gamma_{\nu)}\psi\right] - \\ - g_{\mu\nu}K &= K_Y Y u_{2\mu}u_{2\nu} - Kg_{\mu\nu} = (\rho_f + p_f)u_{2\mu}u_{2\nu} - p_f g_{\mu\nu}. \end{aligned} \tag{2.41}$$

For the FRW metric (2.7), the equations of the f-essence become [13]

$$3H^2 - \rho_f = 0, \tag{2.42}$$

$$2\dot{H} + 3H^2 + p_f = 0, \tag{2.43}$$

$$K_{2Y}\dot{\psi} + 0.5(3HK_{2Y} + \dot{K}_{2Y})\psi - i\gamma^0 K_{2\bar{\psi}} = 0, \tag{2.44}$$

$$K_{2Y}\dot{\bar{\psi}} + 0.5(3HK_{2Y} + \dot{K}_{2Y})\bar{\psi} + iK_{2\psi}\gamma^0 = 0, \tag{2.45}$$

$$\dot{\rho}_f + 3H(\rho_f + p_f) = 0, \tag{2.46}$$

where

$$\rho_f = YK_{2Y} - K_2, \quad p_f = K_2. \tag{2.47}$$

Some properties of f-essence were studied in [52]-[53].

3 Model

In this section, we consider the action (2.1) with the following particular g-essence Lagrangian:

$$K=\alpha_1X+\alpha_2X^n+\alpha_3V_1(\phi)+\beta_1Y+\beta_2Y^m+\beta_3V_2(\bar\psi,\psi)+\eta U_1(\phi)U_2(\bar\psi,\psi), \tag{3.1}$$

where α_j,β_j,η are some real constants. As the search for exact solutions of the coupled system of differential equations (2.9)–(2.14) for the g-essence Lagrangian (3.1) is a very hard job, let us simplify the problem e.g. as $V_2=V_2(u),\beta_2=0,U_1=\phi,U_2=u$. Then the system (2.9)–(2.14) takes the form

$$\begin{aligned}
3H^2-\rho &= 0, \qquad (3.2)\\
2\dot H+3H^2+p &= 0, \qquad (3.3)\\
\ddot\phi+[3H+(\ln(\alpha_1+\alpha_2nX^{n-1}))_t]\dot\phi-\frac{\alpha_3V_{1\phi}+\eta u}{\alpha_1+\alpha_2nX^{n-1}} &= 0, \qquad (3.4)\\
\dot\psi+1.5H\psi-i\beta_1^{-1}\gamma^0(\beta_3V_{2u}\psi+\eta\phi\psi) &= 0, \qquad (3.5)\\
\dot{\bar\psi}+1.5H\bar\psi+i\beta_1^{-1}(\beta_3V_{2u}\bar\psi+\eta\phi\bar\psi)\gamma^0 &= 0, \qquad (3.6)\\
\dot\rho+3H(\rho+p) &= 0, \qquad (3.7)
\end{aligned}$$

where

$$\begin{aligned}
\rho &= \alpha_1X+\alpha_2(2n-1)X^n-\alpha_3V_1-\beta_3V_2-\eta\phi u, \qquad (3.8)\\
p &= \alpha_1X+\alpha_2X^n+\alpha_3V_1+\beta_1Y+\beta_3V_2+\eta\phi u. \qquad (3.9)
\end{aligned}$$

Hence and from (2.23)–(2.24) we get

$$\begin{aligned}
\rho+p &= 2\alpha_1X+2\alpha_2nX^n+\beta_1Y=-2\dot H, \qquad (3.10)\\
\beta_1Y &= -(\beta_3V_2^{'}+\eta\phi)u, \qquad (3.11)\\
u &= \frac{c}{\beta_1a^3}. \qquad (3.12)
\end{aligned}$$

4 Solution

In this section, we want to present the exact solution of the system (3.2)–(3.7). But first note that for six unknown functions $a,\phi,\psi,\bar\psi,V_1,V_2$ we have five differential equations (3.2)–(3.6) so that we need one more equation (see e.g. [47]–[49]). Such an equation we take to be

$$a=\bar a_0\phi^k, \tag{4.1}$$

where $\bar a_0=a_0\phi_0^{-k},\quad k=\lambda/\delta$. Then we obtain the following solution

$$\begin{aligned}
a &= a_0t^\lambda, \qquad (4.2)\\
\phi &= \phi_0t^\delta, \qquad (4.3)\\
\psi_l &= \frac{c_l}{a_0^{1.5}t^{1.5\lambda}}e^{-iD},\quad (l=1,2), \qquad (4.4)\\
\psi_k &= \frac{c_k}{a_0^{1.5}t^{1.5\lambda}}e^{iD},\quad (k=3,4), \qquad (4.5)
\end{aligned}$$

where c_j obey the condition

$$c=|c_1|^2+|c_2|^2-|c_3|^2-|c_4|^2 \tag{4.6}$$

and

$$D=\frac{2}{\beta_1u_0[2(\delta-1)+3\lambda+1]}t^{2(\delta-1)+3\lambda+1}+$$
$$\frac{2^{1-n}\alpha_2n\delta^{2n}\phi_0^{2n}}{u_0[2n(\delta-1)+3\lambda+1]}t^{2n(\delta-1)+3\lambda+1}-\frac{2\lambda}{u_0(3\lambda-1)}t^{3\lambda-1}+D_0. \tag{4.7}$$

This solution is correct if

$$\delta = 2 - 3\lambda \tag{4.8}$$

or

$$\delta = \frac{3\lambda - 2n}{1 - 2n}. \tag{4.9}$$

The corresponding potentials take the form

$$\begin{aligned} V_1(\phi) &= l_1 \left(\frac{\phi}{\phi_0}\right)^{\frac{2(\delta-1)}{\delta}} + l_2 \left(\frac{\phi}{\phi_0}\right)^{\frac{2n(\delta-1)}{\delta}} + l_3 \left(\frac{\phi}{\phi_0}\right)^{\frac{\delta-3\lambda}{\delta}} + V_{10}, \qquad (4.10)\\ V_2(u) &= q_1 \left(\frac{u}{u_0}\right)^{\frac{2(1-\delta)}{3\lambda}} + q_2 \left(\frac{u}{u_0}\right)^{\frac{2n(1-\delta)}{3\lambda}} + q_3 \left(\frac{u}{u_0}\right)^{\frac{3\lambda-\delta}{3\lambda}} + q_4 \left(\frac{u}{u_0}\right)^{\frac{2}{3\lambda}} - \alpha_3\beta_3^{-1}V_{10}. \qquad (4.11) \end{aligned}$$

Here $V_{10} = const, \quad u_0 = ca_0^{-3}\beta_1^{-1}$,

$$\begin{aligned} l_1 &= \frac{\alpha_1\delta^2\phi_0^2(\delta-1+3\lambda)}{2\alpha_3(\delta-1)}, \qquad (4.12)\\ l_2 &= \frac{\alpha_2\delta^{2n}\phi_0^{2n}[(2n-1)(\delta-1)+3\lambda]}{2^n\alpha_3(\delta-1)}, \qquad (4.13)\\ l_3 &= -\frac{\eta\delta\phi_0 u_0}{\alpha_3(\delta-3\lambda)} \qquad (4.14) \end{aligned}$$

and

$$\begin{aligned} q_1 &= 0.5\alpha_1\delta^2\phi_0^2 - \alpha_3 l_1, \qquad (4.15)\\ q_2 &= \alpha_2(2n-1)2^{-n}\delta^{2n}\phi_0^{2n} - \alpha_3 l_2, \qquad (4.16)\\ q_3 &= -\alpha_3 l_3 - \eta\phi_0 u_0, \qquad (4.17)\\ q_4 &= -3\lambda^2. \qquad (4.18) \end{aligned}$$

5 Unified scalar–fermionic Chaplygin gas model of dark energy and dark matter from g-essence with Yukawa interactions

The most popular models of dark energy and dark matter such as e.g. the ΛCDM and a quintesse-CDM model assume that dark energy and dark matter are distinct entities. Another interpretation of the observational data is that dark energy and dark matter are different manifestations of a common structure. The first definite model of this type was proposed in [51], based upon the Chaplygin gas, an exotic perfect fluid obeying the equation of state (EoS)

$$p = -\frac{A}{\rho}, \tag{5.1}$$

which has been extensively studied for its mathematical properties [50]. The general class of models, in which a unification of dark energy and dark matter is achieved through a single entity, is often referred to as quartessence. Among other scenarios of unification that have recently been suggested, interesting attempts are based on k-essence. In this section we extend this scenario to the g-essence model with Yukawa interactions which gives us the Chaplygin gas unified model of dark energy and dark matter. To do it, let us consider the g-essence model given by the following scalar–fermionic DBI Lagrangian:

$$K = U\sqrt{1 + V_1 X + V_2 Y^2}, \tag{5.2}$$

where in general $U = U(\phi, \bar{\psi}, \psi), \quad V_1 = V_1(\phi, \bar{\psi}, \psi), \quad V_2 = V_2(\phi, \bar{\psi}, \psi)$. Note that the g-essence model (5.2) is constrained in two particular cases: i) the scalar DBI model as $U = U(\phi), \quad V_1 =$

$V_1(\phi), \quad V_2 = 0$; ii) the fermionic DBI model as $U = U(\bar{\psi}, \psi), \quad V_1 = 0, \quad V_2 = V_2(\bar{\psi}, \psi)$. Substituting the expression (5.2) into (2.16) we get

$$\begin{aligned} p &= U\sqrt{1 + V_1 X + V_2 Y^2}, &\quad (5.3)\\ \rho &= -\frac{U}{\sqrt{1 + V_1 X + V_2 Y^2}}, &\quad (5.4) \end{aligned}$$

where we assume that

$$U = \eta\phi u, \quad V_1 = V_1(\phi, \bar{\psi}, \psi), \quad V_2 = V_2(\phi, \bar{\psi}, \psi). \tag{5.5}$$

It is the g-essence model with Yukawa interactions $U = \eta\phi u$. These equations give the following EoS:

$$p = -\frac{U^2}{\rho}. \tag{5.6}$$

It is the Chaplygin gas model [51] but with the variable function U (Yukawa interactions). From (5.6) and (2.14), we get

$$\rho = a^{-3}\left[6\int U^2(a)a^5 da + C\right]^{0.5} = z^{-0.5}\left[C + \int U^2(z)dz\right]^{0.5}, \tag{5.7}$$

where $C = const, \quad z = a^6$. From these formulas we obtain the following expression for the EoS parameter:

$$\omega = -\frac{U^2}{\rho^2} = -1 - \frac{zd\ln\rho^2}{dz} = -1 - \frac{d\ln\rho^2}{d\ln z} = -\frac{zU^2}{C + \int U^2(z)dz} = -\frac{zh^{'}}{C + h} = -z[\ln(C+h)]^{'}, \tag{5.8}$$

where $h = \int U^2(z)dz, \quad h^{'} = dh/dz$. In principle, now it is not difficult to construct solutions of the g-essence equations corresponding to the different expressions for U. Here we just present the expressions for the energy density and pressure. Consider some examples.

i) Let $U = \mu = const$. Then from (5.7) and (5.6) we obtain the expressions for the energy density and the pressure

$$\rho = z^{-0.5}\left[C + \mu^2 z\right]^{0.5} = \left[Ca^{-6} + \mu^2\right]^{0.5} \tag{5.9}$$

and

$$p = -\frac{\mu^2 z^{2\nu+0.5}}{\left(C + \frac{\mu^2}{2\nu+1}z^{2\nu+1}\right)^{0.5}} = -\frac{\mu^2}{(Ca^{-6} + \mu^2)^{0.5}}, \tag{5.10}$$

respectively. The EoS parameter is given by

$$\omega = -\frac{\mu^2 z^{2\nu+1}}{C + \frac{\mu^2}{2\nu+1}z^{2\nu+1}}. \tag{5.11}$$

So this example corresponds to the usual Chaplygin gas [51]. As is well known, in this case, for small a ($a^6 << C\mu^{-0.5}$), the energy density and the pressure take the forms, approximately,

$$\rho \approx C^{0.5}a^{-3}, \quad p \approx 0 \tag{5.12}$$

with $\omega = 0$, which corresponds to a matter-dominated universe. For a large value a, it follows that

$$\rho \approx \mu, \quad p \approx -\mu \tag{5.13}$$

that is, $\omega = -1$, which corresponds to a dark energy-dominated universe. So this simple and elegant model smoothly interpolates between a dust dominated phase, where $\rho \approx C^{0.5}a^{-3}$, and a de Sitter phase, where $p \approx -\rho$, through an intermediate regime described by the EoS for stiff matter, $p = \rho$.

ii) Now let us consider the case when $U = \mu z^{\nu}$, where μ and ν are some real constants. In this case, Eq.(5.7) gives the following expression for the energy density:

$$\rho = z^{-0.5}\left[C + \frac{\mu^2}{2\nu+1} z^{2\nu+1}\right]^{0.5} = \left[Ca^{-6} + \frac{\mu^2}{2\nu+1} a^{12\nu}\right]^{0.5}. \tag{5.14}$$

Here ν must be negative, because otherwise, $a \to \infty$ implies $\rho \to \infty$, which is not the case for expanding Universe. The equation (5.6) gives the expression for the pressure,

$$p = -\frac{\mu^2 z^{2\nu+0.5}}{\left(C + \frac{\mu^2}{2\nu+1} z^{2\nu+1}\right)^{0.5}} = -\frac{\mu^2 a^{12\nu}}{\left(Ca^{-6} + \frac{\mu^2}{2\nu+1} a^{12\nu}\right)^{0.5}}. \tag{5.15}$$

The EoS parameter is given by

$$\omega = -\frac{\mu^2 z^{2\nu+1}}{C + \frac{\mu^2}{2\nu+1} z^{2\nu+1}} = -\frac{\mu^2 a^{12\nu}}{Ca^{-6} + \frac{\mu^2}{2\nu+1} a^{12\nu}} = -\frac{\mu^2}{Ca^{-6(1+2\nu)} + \frac{\mu^2}{2\nu+1}}. \tag{5.16}$$

The deceleration parameter q has the expression

$$q = -\frac{\ddot{a}}{aH^2} = \frac{\rho + 3p}{2\rho} = \frac{Ca^{-6} + \frac{2\mu^2(3\nu+2)}{2\nu+1} a^{12\nu}}{2(Ca^{-6} + \frac{\mu^2}{2\nu+1} a^{12\nu})}. \tag{5.17}$$

For accelerating universe, q must be negative i.e., $\ddot{a} > 0$. Hence we have

$$a^{-6(2\nu+1)} < -\frac{2\mu^2(3\nu+2)}{C(2\nu+1)}. \tag{5.18}$$

This means that for a small value of the scale factor we have a decelerating universe while for large values of scale factor we have an accelerating universe. The transition between these two phases occurs when the scale factor is equal to

$$a = a_c = -\left[\frac{2\mu^2(3\nu+2)}{C(2\nu+1)}\right]^{\frac{1}{-6(2\nu+1)}}. \tag{5.19}$$

6 Unified scalar–fermionic generalized Chaplygin gas model of dark energy and dark matter from g-essence with Yukawa interactions

In the previous section we constructed the unified scalar–fermionic Chaplygin gas model of dark energy and dark matter using the particular g-essence with Yukawa interactions having the scalar–fermionic DBI Lagrangian form (5.2). In this section we extend these results of the previous section for the scalar–fermionic generalized Chaplygin gas case. For this purpose, we consider the particular g-essence model with Yukawa interactions which has the following scalar–fermionic DBI Lagrangian form

$$K = U(1 + V_1 X^{\frac{1}{2n}} + V_2 Y^{\frac{1}{n}})^n, \tag{6.1}$$

where

$$U = \eta\phi u, \quad V_1 = V_1(\phi, \bar{\psi}, \psi), \quad V_2 = V_2(\phi, \bar{\psi}, \psi). \tag{6.2}$$

Substituting this expression into (2.16) we get

$$p = U(1 + V_1 X^{\frac{1}{2n}} + V_2 Y^{\frac{1}{n}})^n, \tag{6.3}$$

$$\rho = -U(1 + V_1 X^{\frac{1}{2n}} + V_2 Y^{\frac{1}{n}})^{n-1}. \tag{6.4}$$

These equations give

$$p = -\frac{(-U)^{\frac{1}{1-n}}}{\rho^{\frac{n}{1-n}}} = -\frac{(-U)^{\alpha+1}}{\rho^{\alpha}} = -\frac{A}{\rho^{\alpha}}, \tag{6.5}$$

where $\alpha=n(1-n)^{-1}$, $A=(-U)^{\alpha+1}$. It is the scalar–fermionic generalized Chaplygin gas model [51]. From (2.16) and (6.5) we get

$$\rho=a^{-3}\left[3(1+\alpha)\int(-U)^{\alpha+1}a^{3(1+\alpha)-1}da+C\right]^{\frac{1}{1+\alpha}}=z^{-\frac{1}{1+\alpha}}\left[C+\int(-U)^{\alpha+1}dz\right]^{\frac{1}{1+\alpha}}, \tag{6.6}$$

where $C=const$, $z=a^{3(1+\alpha)}$. At the same time, for the pressure we obtain the following expression

$$p=-(-U)^{\alpha+1}z^{\frac{\alpha}{1+\alpha}}\left[C+\int(-U)^{\alpha+1}dz\right]^{-\frac{\alpha}{1+\alpha}}. \tag{6.7}$$

These formulas give the following expression for the EoS parameter:

$$\omega=-\frac{z(-U)^{\alpha+1}}{C+\int(-U)^{\alpha+1}dz}. \tag{6.8}$$

Now we can consider different types solutions of the g-essence equations. For example, if for simplicity, we consider the case $\omega=const$ then from (6.8) we obtain the following expression for the function U:

$$U=-A_0^{\frac{1}{1+\alpha}}a^{-3(1+\omega)}, \tag{6.9}$$

where A_0 is an integration constant. Let us consider some examples: i) if $\omega=-1$ (the de Sitter case) then $U=-A_0^{\frac{1}{1+\alpha}}$; ii) if $\omega=0$ (the dust case) then $U=-A_0^{\frac{1}{1+\alpha}}a^{-3}$; iii) if $\omega=1$ (the stiff matter case) then $U=-A_0^{\frac{1}{1+\alpha}}a^{-6}$ and so on. Now the construction of solutions of the Friedmann, Klein–Gordon and Dirac equations is a formal problem so we omit it here.

7 Scalar–fermionic DBI generalization of the g-essence model with Yukawa interaction (3.1)

Our aim in this section is to construct the scalar–fermionic DBI generalization of the g-essence model with Yukawa interaction (3.1). To do it, we note that the model (3.1) is some approximation of the following scalar–fermionic DBI model:

$$K=\epsilon A\left\{\sqrt{1+A^{-1}[2\alpha_1X+2\alpha_2X^n+2\beta_1Y+2\beta_2Y^m]}-1\right\}+V, \tag{7.1}$$

where $\epsilon=+1$, $A=const$ and

$$V=\alpha_3V_1(\phi)+\beta_3V_2(\bar{\psi},\psi)+\eta U_1(\phi)U_2(\bar{\psi},\psi). \tag{7.2}$$

From (7.1) we get

$$\begin{aligned} p &= \epsilon A\left\{\sqrt{1+A^{-1}[2\alpha_1X+2\alpha_2X^n+2\beta_1Y+2\beta_2Y^m]}-1\right\}+V, && (7.3)\\ \rho &= \frac{2\epsilon\alpha_2(n-1)X^n-\epsilon\beta_1Y+\epsilon\beta_2(m-2)Y^m-\epsilon A}{\sqrt{1+A^{-1}[2\alpha_1X+2\alpha_2X^n+2\beta_1Y+2\beta_2Y^m]}}+\epsilon A-V. && (7.4)\end{aligned}$$

The study the system of equations of g-essence (2.9)–(2.14) with the expressions for the pressure and the energy density given by (7.3)–(7.4) is a very hard job. Let us simplify the problem. Let $\alpha_2=\beta_1=0$ and $m=2$. Then equations (7.3)–(7.4) take the form

$$\begin{aligned} p &= \epsilon A\left\{\sqrt{1+A^{-1}[2\alpha_1X+2\beta_2Y^2]}-1\right\}+V, && (7.5)\\ \rho &= -\frac{\epsilon A}{\sqrt{1+A^{-1}[2\alpha_1X+2\beta_2Y^2]}}+\epsilon A-V && (7.6)\end{aligned}$$

which corresponds to the EoS

$$p=-\frac{A^2}{\rho-\epsilon A+V}-\epsilon A+V. \tag{7.7}$$

The corresponding EoS parameter is given by

$$\omega = -\frac{A^2}{\rho(\rho - \epsilon A + V)} - \frac{\epsilon A - V}{\rho} = \frac{A^2}{\epsilon A - V}\left(\frac{1}{\rho} - \frac{1}{\rho - \epsilon A + V}\right). \tag{7.8}$$

Now let us consider two particular limit cases when $\rho >> \epsilon A - V$ or $\rho << \epsilon A - V$.

i) First we consider the case when $\rho >> \epsilon A - V$. Let us rewrite the equation (5.7) as

$$p = -\frac{A^2}{\rho}\left(\frac{1}{1 - \frac{\epsilon A - V}{\rho}}\right) - \epsilon A + V. \tag{7.9}$$

Hence, in the first approximation, we get

$$p = -\frac{A^2}{\rho} - \epsilon A + V \tag{7.10}$$

which corresponds to the Chaplygin gas case [51]. Now let us consider the second approximation. Then from (7.9) we obtain

$$p = -\frac{A^2}{\rho - \epsilon A + V} \approx -\frac{A^2}{\rho}\left(1 + \frac{\epsilon A - V}{\rho}\right) - \epsilon A + V = -\frac{A^2}{\rho} - \frac{A^2(\epsilon A - V)}{\rho^2} - \epsilon A + V. \tag{7.11}$$

ii) Now let us consider the second case when $\rho << \epsilon A - V$. Then from (7.7) we get

$$p = -\frac{A^2}{\rho - \epsilon A + V} \approx \frac{A^2}{\epsilon A - V}\left(1 + \frac{\rho}{\epsilon A - V}\right) - \epsilon A + V = \Lambda_g + \frac{A^2}{(\epsilon A - V)^2}\rho, \tag{7.12}$$

where

$$\Lambda_g = \frac{A^2}{\epsilon A - V} - \epsilon A + V. \tag{7.13}$$

In this case, the EoS parameter is given by

$$\omega = \frac{A^2}{(\epsilon A - V)^2} + \frac{\Lambda_g}{\rho}. \tag{7.14}$$

8 Conclusion

Recently, the so-called g-essence has been proposed to be a candidate of dark energy. It has the non-standard kinetic scalar and fermionic terms and has interesting properties. We expect that g-essence has rich properties and when it is used in cosmology, its rich properties could have some interesting consequences. In this work, we found the exact power law solution of the particular g-essence model with the Yukawa scalar–fermionic interactions. The corresponding scalar and fermionic potentials are presented. Of course, there might be other exact solutions of the g-essence different from the power law solution. Anyway, our results obtained in the present work showed that the g-essence with the Yukawa interactions can describes the accelerated expansions of the universe. In fact, we have shown that some particular g-essence models with Yukawa interactions correspond to the usual and generalized Chaplygin gas unified models of dark energy and dark matter. Also we presented some scalar–fermionic DBI models which again can describe the unified dark energy–dark matter system.

9 Acknowledgement

We would like to thank the anonymous referees for providing us with constructive comments and suggestions to improve this work.

References

[1] Perlmutter S. et al. *Measurements of omega and lambda from 42 high-redshift supernovae*, Astrophys. J. **517**, 565-586 (1999) [astro-ph/9812133].

[2] Riess et al. *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, Astron. J. **116**, 1009-1038 (1998) [astro-ph/9805201].

[3] Armendariz-Picon C., Damour T., Mukhanov V.F. *k-inflation*, Phys. Lett. **B458**, 209-218 (1999) [hep-th/9904075].

[4] Armendariz-Picon C., Mukhanov V.F., Steinhardt P.J. *Essentials of k-essence*, Phys. Rev. **D63**, 103510 (2001) [astro-ph/0006373].

[5] Armendariz-Picon C., Mukhanov V.F., Steinhardt P.J. *A dynamical solution to the problem of a small cosmological constant and late-time cosmic acceleration*, Phys. Rev. Lett. **85**, 4438-4441 (2000) [astro-ph/0004134].

[6] Chiba T., Okabe T., Yamaguchi M. *Kinetically driven quintessence*, Phys. Rev. **D62**, 023511 (2000) [astro-ph/9912463].

[7] De Putter R., Linder E.V. *Kinetic k-essence and Quintessence*, Astropart. Phys. **28**, 263-272 (2007), [arXiv:0705.0400]

[8] Myrzakulov R. *F(T) gravity and k-essence*, [arXiv:1008.4486]

[9] Myrzakulov R. *Pure kinetic k-essence as the cosmic speed-up*, IJTP, **50**, 1876-1886 (2011) [arXiv:1008.0779]

[10] Ribas M.O., Devecchi F.P., Kremer G.M. *Fermions as sources of accelerated regimes in cosmology*, Phys. Rev.**D72**, 123502 (2005) [arXiv:gr-qc/0511099]

[11] Samojeden L.L., Devecchi F.P., Kremer G.M. *Fermions in Brans-Dicke cosmology*, Phys. Rev. **D81**, 027301 (2010) [arXiv:1001.2285].

[12] Samojeden L.L., Kremer G.M., Devecchi F.P. *Accelerated expansion in bosonic and fermionic 2D cosmologies with quantum effects*, Europhys. Lett. **87**, 10001 (2009) [arXiv:0906.3315].

[13] Myrzakulov R. *Fermionic k-essence*, [arXiv:1011.4337v5]

[14] Esmakhanova K., Myrzakulov N., Nugmanova G., Chechin L., Myrzakulov R. *Integrable and nonintegrable FRW cosmological models induced by some second-order ordinary differential equations*, [arXiv:1104.3705]

[15] Nugmanova G.N., Myrzakul Sh.R., Razina O.V., Esmakhanova K.R., Serikbayev N.S., Myrzakulov R. *Some cosmological aspects of Horava-Lifshitz gravity: integrable and nonintegrable models*, [arXiv:1104.5374]

[16] Myrzakul Sh., Esmakhanova K., Myrzakulov K., Nugmanova G., Myrzakulov R. *FRW cosmological models with integrable and nonintegrable differential equations of state*, [arXiv:1105.2771].

[17] Yerzhanov K. K., Tsyba P. Yu., Myrzakul Sh. R., Kulnazarov I. I., Myrzakulov R. *Accelerated expansion of the Universe driven by G-essence*, [arXiv:1012.3031]

[18] Razina O., Tsyba P., Kulnazarov I., Myrzakul Sh., Yerzhanov K., Myrzakulov R. *G-essence cosmologies with Yukawa type interactions*, [arXiv:1101.0729]

[19] Ribas M.O., Kremer G.M. *Fermion fields in Einstein-Cartan theory and the accelerated-decelerated transtion in a primordial Universe*, Grav. Cosmol. **16**, 173-177 (2010) [arXiv:0902.2696].

[20] Cai Y.F., Wang J. *Dark Energy Model with Spinor Matter and Its Quintom Scenario*, Class. Quant. Grav., **25**, 165014 (2008) [arXiv:0806.3890]

[21] Wang J., Cui S.-W., Zhang C.-M. *Thermodynamics of Spinor Quintom*, Phys. Lett., **B683**, 101-107 (2010) [arXiv:0806.3890]

[22] Ribas M.O., Devecchi F.P., Kremer G.M. *Cosmological model with non-minimally coupled fermionic field*, Europhys. Lett. **81**, 19001 (2008) [arXiv:0710.5155]

[23] Rakhi R., Vijayagovindan G.V., Indulekha K. *A cosmological model with fermionic field*, [arXiv:0912.1222]

[24] Rakhi R., Vijayagovindan G.V., Noble P. A., Indulekha K. *A cosmological model with fermionic field and Gauss-Bonnet term*, IJMPA, **25**, 1267-1278 (2010) [arXiv:0910.3761]

[25] Chimento L.P., Devecchi F.P., Forte M., Kremer G.M. *Phantom cosmologies and fermions*, Class.Quant. Grav. **25**, 085007 (2008) [arXiv:0707.4455]

[26] Anischenko S.V., Cherkas S.L., Kalashnikov V.L. *Cosmological Production of Fermions in a Flat Friedmann Universe with Linearly Growing Scale Factor: Exactly Solvable Model*, [arXiv:0911:0769]

[27] Saha B. *Nonlinear spinor field in cosmology*, Phys. Rev. **D69**, 124006 (2004) [arXiv:gr-qc/0308088]

[28] Saha B., Shikin G.N. *Nonlinear Spinor Field in Bianchi type-I Universe filled with Perfect Fluid: Exact Self-consistent Solutions*, J. Math. Phys. **38**, 5305 (1997)

[29] Saha B. *Spinor Field in Bianchi type-I Universe: regular solutions*, Phys. Rev. **D64**, 123501 (2001) [arXiv:gr-qc/0107013]

[30] Saha B. *Spinor felds in Bianchi type-I Universe Physics of Particles and Nuclei*, Physics of Particles and Nuclei, **37**, Suppl., S13 (2006)

[31] Saha B. *Nonlinear spinor field in Bianchi type-I cosmology: Inflation, isotropization, and late time acceleration*, Phys. Rev. **D74**, 124030 (2006)

[32] Vakili B., Sepangi H.R. *Time reparameterization in Bianchi type I spinor cosmology*, Annals Phys. **323**, 548-565 (2008) [arXiv:0709.2988].

[33] Wei H. *Spinor Dark Energy and Cosmological Coincidence Problem*, Phys. Lett. B **695**, 307-311 (2011) [arXiv:1002.4230].

[34] Dereli T., Ozdemir N., Sert O. *Einstein-Cartan-Dirac Theory in (1+2)-Dimensions*, [arXiv:1002.0958]

[35] Balantekin A.B., Dereli T. *An Exact Cosmological Solution of the Coupled Einstein-Majorana Fermion-Scalar Field Equations*, Phys. Rev. **D75**, 024039 (2007), [arXiv:gr-qc/0701025].

[36] Armendariz-Picon C., Greene P. *Spinors, Inflation, and Non-Singular Cyclic Cosmologies*, Gen.Relativ. Gravit. **35**, 1637-1658 (2003), [arXiv:hep-th/0301129]

[37] Weinberg S. *Gravitation and Cosmology* (John Wiley & Sons, New York, 1972), ibid. *Cosmology* (Cambridge, New York, 2007).

[38] Wald R.M. *General Relativity*, (The University of Chicago Press, Chicago, 1984).

[39] Ryder L.H. *Quantum Field Theory* (Cambridge University Press, Cambridge, 1996).

[40] Birrell N.D., Davies P.C.W. *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).

[41] Zanusso O., Zambelli L., Vacca G.P., Percacci R. *Gravitational corrections to Yukawa systems*, [arXiv:0904.0938]

[42] Rodigast A., Schuster T. *Gravitational Corrections to Yukawa and* φ^4 *Interactions*, [arXiv:0908.2422]

[43] Loeb A., Weiner N. *Cores in Dwarf Galaxies from Dark Matter with a Yukawa Potential*, [arXiv:1011.6374]

[44] Miao S.P., Woodard R.P. *Leading Log Solution for Inflationary Yukawa*, [gr-qc/0602110]

[45] Woodard R.P. *Generalizing Starobinskii's Formalism to Yukawa Theory & to Scalar QED*, [gr-qc/0608037]

[46] Atazadeh K., Sepangi H.R. *Accelerated expansion in modified gravity with a Yukawa-like term*, [gr-qc/0602028]

[47] Elizalde E., Myrzakulov R., Obukhov V.V., Saez-Gomez D. *LambdaCDM epoch reconstruction from F(R,G) and modified Gauss-Bonnet gravities*, Class.Quant.Grav., **27**, 095007 (2010) [arXiv:1001.3636]

[48] Myrzakulov R., Saez-Gomez D., Tureanu A. *On the* Λ*CDM Universe in* $f(G)$ *gravity*, Gen. Rel. Grav., **43**, 1671-1684 (2011) [arXiv:1009.0902]

[49] Nojiri S., Odintsov S.D. *Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models*, Physics Reports, (2011), [arXiv:1011.0544]

[50] Jackiw R. *Lectures on fluid dynamics*, Springer-Verlag, New-York, 2002.

[51] Kamenshchik A.Y., Moschella U., Pasquier V. *An alternative to quintessence*, Phys. Lett. B, **511**, 265 (2001)
Bento M.C., Bertolami O., Sen A.A. *Generalized Chaplygin Gas, Accelerated Expansion and Dark Energy-Matter Unification*, Phys. Rev. D, **66**, 043507 (2002)

[52] Tsyba P., Yerzhanov K., Esmakhanova K., Kulnazarov I., Nugmanova G., Myrzakulov R. *Reconstruction of f-essence and fermionic Chaplygin gas models of dark energy*, [arXiv:1103.5918]

[53] Jamil M., Myrzakulov Y., Razina O., Myrzakulov R. *Modified Chaplygin Gas and Solvable F-essence Cosmologies*, [arXiv:1107.1008]